



A counterexample to the Fourteenth Problem of Hilbert in dimension four

Shigeru Kuroda¹

Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606-8502, Japan

Received 13 August 2003

Available online 26 June 2004

Communicated by Paul Roberts

Abstract

This paper presents a family of new counterexamples to Hilbert's Fourteenth Problem. They are realized as subfields of the rational function field of four variables over a field of characteristic zero of transcendence degree three. It was previously not known whether any counterexample could be found as a subfield of a rational function field of four variables, or whether counterexamples with minimal transcendence degree three could exist.

© 2004 Elsevier Inc. All rights reserved.

1. Introduction

Let K be a field of characteristic zero, $K[\mathbf{x}] = K[x_1, \dots, x_n]$ the polynomial ring in n variables over K , and $K(\mathbf{x})$ its field of fractions. Then, the Fourteenth Problem of Hilbert asks whether the K -subalgebra $L \cap K[\mathbf{x}]$ of $K[\mathbf{x}]$ is finitely generated for a subfield L of $K(\mathbf{x})$ containing K . The first counterexample to this problem was found by Nagata [12] in 1958 in the case where $n = 2s^2$ for each integer $s \geq 4$. In 1990, Roberts [14] constructed a new counterexample when $n = 7$. Kojima and Miyanishi [6] gave a similar counterexample for each odd number $n \geq 7$. Recently, we generalized Roberts' example further in [8], and showed a detailed sufficient condition for certain invariant subrings of $K[\mathbf{x}]$ not to be

E-mail address: kuroda@kurims.kyoto-u.ac.jp.

¹ Partly supported by the Grant-in-Aid for JSPS Fellows, The Ministry of Education, Science, Sports and Culture, Japan.

finitely generated in the case where $n \geq 7$. In lower dimensions, Freudenburg [4] gave a counterexample for $n = 6$, and Daigle and Freudenburg [1] gave one for $n = 5$.

While there exists a counterexample for each $n \geq 5$, the answer to the Fourteenth Problem of Hilbert is known to be affirmative for $n \leq 2$. Actually, Zariski [15] showed that $L \cap K[\mathbf{x}]$ is finitely generated if the transcendence degree of L over K is at most two. In case of $n = 3, 4$, there were only positive solutions for some special cases (cf. [2, 7, 9, 11]).

In the case of Nagata's counterexample, the transcendence degree of L over K is four. After giving the counterexample, Nagata [13] proposed two open problems concerning the Fourteenth Problem of Hilbert. His second problem asks the answer to the Fourteenth Problem in the case where the transcendence degree of L over K is three.

In the present paper, we give the first counterexample to the Fourteenth Problem of Hilbert for $n = 4$. We also settle Nagata's second problem negatively by our new counterexample. Furthermore, we show that the counterexample cannot be obtained as the kernel of any locally nilpotent derivation. Here, a K -linear map $D: A \rightarrow A$ of a commutative K -algebra A is called a *derivation* if $D(fg) = D(f)g + fD(g)$ for any $f, g \in A$. Then, the kernel

$$A^D = \{f \in A \mid D(f) = 0\}$$

of D is a K -subalgebra of A . A derivation D on A is said to be *locally nilpotent* if, for each $f \in A$, there exists $r > 0$ such that $D^r(f) = 0$. We note that each of the counterexamples in [1, 4, 6, 8, 14] is given as the kernel of certain locally nilpotent derivation on $K[\mathbf{x}]$.

Assume that $n = 4$. Let γ and $\delta_{i,j}$ be integers for $1 \leq i \leq 3$ and $1 \leq j \leq 4$ such that $\gamma, \delta_{i,j} \geq 1$ and $\delta_{i,4} \geq 0$ for $1 \leq i, j \leq 3$, and let $K(\Pi)$ be the subfield of $K(\mathbf{x})$ generated by

$$\begin{aligned} \Pi_1 &= x_4^\gamma - x_1^{-\delta_{1,1}} x_2^{\delta_{1,2}} x_3^{\delta_{1,3}} x_4^{\delta_{1,4}}, & \Pi_2 &= x_4^\gamma - x_1^{\delta_{2,1}} x_2^{-\delta_{2,2}} x_3^{\delta_{2,3}} x_4^{\delta_{2,4}}, \\ \Pi_3 &= x_4^\gamma - x_1^{\delta_{3,1}} x_2^{\delta_{3,2}} x_3^{-\delta_{3,3}} x_4^{\delta_{3,4}} \end{aligned} \quad (1.1)$$

over K .

The following is the main result of this paper.

Theorem 1.1. *Assume that $n = 4$. If*

$$\frac{\delta_{1,1}}{\delta_{1,1} + \min\{\delta_{2,1}, \delta_{3,1}\}} + \frac{\delta_{2,2}}{\delta_{2,2} + \min\{\delta_{3,2}, \delta_{1,2}\}} + \frac{\delta_{3,3}}{\delta_{3,3} + \min\{\delta_{1,3}, \delta_{2,3}\}} < 1, \quad (1.2)$$

then $K(\Pi) \cap K[\mathbf{x}]$ is not finitely generated over K . Moreover, $K(\Pi) \cap K[\mathbf{x}]$ is not contained in the kernel $K[\mathbf{x}]^D$ of any nonzero locally nilpotent derivation D on $K[\mathbf{x}]$.

Clearly, the transcendence degree of $K(\Pi)$ over K is at most three. Hence, Theorem 1.1 implies that Nagata's second problem has a negative answer.

We remark that $K(\Pi) \cap K[\mathbf{x}]$ is not equal to the invariant subring $K[\mathbf{x}]^G$ of $K[\mathbf{x}]$ for $n = 4$ for any algebraic action of an algebraic group G by the following reason, which was

pointed out by the referee. Without loss of generality, we may assume that G acts on $K[\mathbf{x}]$ faithfully. Suppose to the contrary that $K(\Pi) \cap K[\mathbf{x}] = K[\mathbf{x}]^G$. Then, G is not reductive by the Hilbert finiteness theorem. Hence, G contains the one-dimensional additive group G_a as a subgroup, and so $K(\Pi) \cap K[\mathbf{x}] \subset K[\mathbf{x}]^{G_a}$. This contradicts Theorem 1.1, since $K[\mathbf{x}]^{G_a}$ is equal to the kernel $K[\mathbf{x}]^D$ of a nonzero locally nilpotent derivation D on $K[\mathbf{x}]$.

Note: the author recently constructed a counterexample for $n = 3$ [10]. Hence, Hilbert's Fourteenth Problem is settled for every n at last.

2. The structure of $K(\Pi) \cap K[\mathbf{x}]$

Let $K[\mathbf{y}]$ and $K[\mathbf{y}, \mathbf{y}^{-1}]$ be the polynomial ring and the Laurent polynomial ring in y_1, y_2, y_3, y_4 over K , respectively, and $K(\mathbf{y})$ the field of fractions of $K[\mathbf{y}]$. For $a = (a_1, a_2, a_3, a_4) \in \mathbf{Z}^4$, we denote $\mathbf{x}^a = x_1^{a_1} x_2^{a_2} x_3^{a_3} x_4^{a_4}$ and $\mathbf{y}^a = y_1^{a_1} y_2^{a_2} y_3^{a_3} y_4^{a_4}$. If f is the Laurent polynomial $\sum_{a \in \mathbf{Z}^4} \lambda_a \mathbf{x}^a$ or $\sum_{a \in \mathbf{Z}^4} \lambda_a \mathbf{y}^a$, then define the *support* $\text{supp}(f)$ of f by

$$\text{supp}(f) = \{a \in \mathbf{Z}^4 \mid \lambda_a \neq 0\}.$$

We denote by $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4$ the coordinate unit vectors of \mathbf{R}^4 , and by $\mathbf{Z}_{\geq 0}$ the set of nonnegative integers.

Let $K[\mathbf{x}]' = K[\mathbf{x}^{\pm \delta_1}, \mathbf{x}^{\pm \delta_2}, \mathbf{x}^{\pm \delta_3}, x_4^{\pm \gamma}]$, and $K(\mathbf{x})'$ its field of fractions, where

$$\begin{aligned} \delta_1 &= (-\delta_{1,1}, \delta_{1,2}, \delta_{1,3}, \delta_{1,4}), & \delta_2 &= (\delta_{2,1}, -\delta_{2,2}, \delta_{2,3}, \delta_{2,4}), \\ \delta_3 &= (\delta_{3,1}, \delta_{3,2}, -\delta_{3,3}, \delta_{3,4}) \end{aligned} \quad (2.1)$$

and $\delta_{i,j}$ and γ are as defined above. Then, we may define an isomorphism $\Phi: K[\mathbf{y}, \mathbf{y}^{-1}] \rightarrow K[\mathbf{x}]'$ of K -algebras by $y_i \mapsto \mathbf{x}^{\delta_i}$ for $i = 1, 2, 3$ and $y_4 \mapsto x_4^\gamma$. To see this, it suffices to verify that $\delta_1, \delta_2, \delta_3$ and $\gamma \mathbf{e}_4$ are linearly independent over \mathbf{R} . Suppose the contrary. Then, there exists $(\lambda_1, \lambda_2, \lambda_3) \in \mathbf{R}^3 \setminus \{0\}$ such that the first three components of $\sum_{i=1}^3 \lambda_i \delta_i$ are zero. It implies that $\lambda_1, \lambda_2, \lambda_3 > 0$ or $\lambda_1, \lambda_2, \lambda_3 < 0$. So, without loss of generality, we may assume that $0 < \lambda_1 \leq \lambda_2 \leq \lambda_3$. Since $\delta_{1,1} = (\lambda_2/\lambda_1)\delta_{2,1} + (\lambda_3/\lambda_1)\delta_{3,1}$, we have $\delta_{1,1} \geq 2 \min\{\delta_{2,1}, \delta_{3,1}\}$. Hence, $\xi_1 \geq 2/3$. Here, we set

$$\begin{aligned} \xi_1 &= \frac{\delta_{1,1}}{\delta_{1,1} + \min\{\delta_{2,1}, \delta_{3,1}\}}, & \xi_2 &= \frac{\delta_{2,2}}{\delta_{2,2} + \min\{\delta_{3,2}, \delta_{1,2}\}}, \\ \xi_3 &= \frac{\delta_{3,3}}{\delta_{3,3} + \min\{\delta_{1,3}, \delta_{2,3}\}}. \end{aligned} \quad (2.2)$$

Since $\delta_{2,2} = (\lambda_1/\lambda_2)\delta_{1,2} + (\lambda_3/\lambda_2)\delta_{3,2}$, we have $\delta_{2,2} > \min\{\delta_{3,2}, \delta_{1,2}\}$. Hence, $\xi_2 > 1/2$. This contradicts (1.2). Therefore, Φ is an isomorphism, and hence it induces an isomorphism $K(\mathbf{y}) \rightarrow K(\mathbf{x})'$.

Consider the derivation E on $K[\mathbf{y}]$ defined by $E(y_i) = 1$ for each i . It is naturally extended to derivations on $K[\mathbf{y}, \mathbf{y}^{-1}]$ and $K(\mathbf{y})$, respectively. Since $E(y_4 - y_i) = 0$ and $\Pi_i = \Phi(y_4 - y_i)$ for $i = 1, 2, 3$, we have $K(\Pi) \subset \Phi(K(\mathbf{y})^E)$.

Lemma 2.1. *It follows that $K[\mathbf{y}, \mathbf{y}^{-1}]^E = K[\mathbf{y}]^E = K[y_4 - y_1, y_4 - y_2, y_4 - y_3]$.*

Proof. Consider the automorphism $\sigma : K[\mathbf{y}] \rightarrow K[\mathbf{y}]$ defined by $y_i \mapsto y_4 - y_i$ for $i = 1, 2, 3$ and $y_4 \mapsto y_4$. Then, $\sigma^{-1} \circ E \circ \sigma$ is equal to the partial derivative $\partial/\partial y_4$. Hence,

$$K[\mathbf{y}]^E = \sigma(K[\mathbf{y}]^{\sigma^{-1} \circ E \circ \sigma}) = \sigma(K[y_1, y_2, y_3]) = K[y_4 - y_1, y_4 - y_2, y_4 - y_3].$$

Now, we show $K[\mathbf{y}, \mathbf{y}^{-1}]^E = K[\mathbf{y}]^E$. It is clear that $K[\mathbf{y}, \mathbf{y}^{-1}]^E \supset K[\mathbf{y}]^E$. For the converse, suppose that $E(f/\mathbf{y}^a) = 0$ for some $f \in K[\mathbf{y}] \setminus \{0\}$ and $a \in (\mathbb{Z}_{\geq 0})^4 \setminus \{0\}$ such that $\gcd(f, \mathbf{y}^a) = 1$. Then,

$$E(f) = E((f/\mathbf{y}^a)\mathbf{y}^a) = E(f/\mathbf{y}^a)\mathbf{y}^a + (f/\mathbf{y}^a)E(\mathbf{y}^a) = (f/\mathbf{y}^a)E(\mathbf{y}^a),$$

which is in $K[\mathbf{y}]$. Hence, $E(\mathbf{y}^a) = h\mathbf{y}^a$ for some $h \in K[\mathbf{y}]$ by the assumption that $\gcd(f, \mathbf{y}^a) = 1$. Note that E is a locally nilpotent derivation on $K[\mathbf{y}]$, and an eigenvalue of a nonzero locally nilpotent derivation must be zero (see [3, Proposition 1.3.32(ii)]). Hence, we have $E(\mathbf{y}^a) = 0$. It implies that $y_i \in K[\mathbf{y}]^E$ for some i , since $K[\mathbf{y}]^E$ is factorially closed in $K[\mathbf{y}]$ (see [3, Proposition 1.3.32(iii)]) and $\mathbf{y}^a \neq 1$. This contradicts the definition of E . Therefore, $K[\mathbf{y}, \mathbf{y}^{-1}]^E \subset K[\mathbf{y}]^E$. \square

Let us denote $K[\Pi] = K[\Pi_1, \Pi_2, \Pi_3]$. By Lemma 2.1, $K[\mathbf{y}]^E$ is isomorphic to $K[\Pi]$ via Φ ; in fact, $K[\Pi] = \Phi(K[\mathbf{y}]^E)$.

Lemma 2.2. *It follows that $K(\Pi) \cap K[\mathbf{x}] = K[\Pi] \cap K[\mathbf{x}]$.*

Proof. It is clear that $K(\Pi) \cap K[\mathbf{x}] \supset K[\Pi] \cap K[\mathbf{x}]$. We show that $K(\Pi) \cap K[\mathbf{x}] \subset K[\mathbf{x}]'$ in the following paragraph. Then, $K(\Pi) \cap K[\mathbf{x}] \subset K[\Pi] \cap K[\mathbf{x}]$ is proved as follows. By Lemma 2.1, we have

$$\begin{aligned} \Phi^{-1}(K(\Pi) \cap K[\mathbf{x}]') &= \Phi^{-1}(K(\Pi)) \cap \Phi^{-1}(K[\mathbf{x}]') \subset K(\mathbf{y})^E \cap K[\mathbf{y}, \mathbf{y}^{-1}] \\ &= K[\mathbf{y}, \mathbf{y}^{-1}]^E = K[\mathbf{y}]^E. \end{aligned}$$

Hence, $K(\Pi) \cap K[\mathbf{x}]' \subset K[\Pi]$, since $\Phi(K[\mathbf{y}]^E) = K[\Pi]$. By the assumption that $K(\Pi) \cap K[\mathbf{x}] \subset K[\mathbf{x}]'$, we have $K(\Pi) \cap K[\mathbf{x}] \subset K[\Pi]$. Thus, we get $K(\Pi) \cap K[\mathbf{x}] \subset K[\Pi] \cap K[\mathbf{x}]$.

Since $K(\Pi) \cap K[\mathbf{x}] \subset K(\mathbf{x})' \cap K[\mathbf{x}, \mathbf{x}^{-1}]$, we show that $K(\mathbf{x})' \cap K[\mathbf{x}, \mathbf{x}^{-1}] \subset K[\mathbf{x}]'$. Suppose that there exist $F, G \in K[\mathbf{x}]'$ such that $F/G \in K[\mathbf{x}, \mathbf{x}^{-1}] \setminus K[\mathbf{x}]'$. If h is the sum of terms of F/G contained in $K[\mathbf{x}]'$, then $F - Gh \in K[\mathbf{x}]'$ and $(F - Gh)/G = F/G - h \in K[\mathbf{x}, \mathbf{x}^{-1}] \setminus K[\mathbf{x}]'$. So, by replacing F with $F - Gh$, we may assume that any monomial appearing in F/G is not in $K[\mathbf{x}]'$. For each $f \in K[\mathbf{x}, \mathbf{x}^{-1}] \setminus \{0\}$, let $\text{in}(f)$ denote the maximal monomial appearing in f with nonzero coefficient for the total ordering \preccurlyeq on the set of monomials defined by $\mathbf{x}^a \preccurlyeq \mathbf{x}^b$ if the first nonzero component of $b - a$ is positive. Then, $\text{in}(F)/\text{in}(G) \in K[\mathbf{x}]'$, since $\text{in}(F), \text{in}(G) \in K[\mathbf{x}]'$. This is a contradiction, since $\text{in}(F)/\text{in}(G) = \text{in}(F/G)$, which is not in $K[\mathbf{x}]'$. Hence, $K(\mathbf{x})' \cap K[\mathbf{x}, \mathbf{x}^{-1}] \subset K[\mathbf{x}]'$. \square

Lemma 2.3. *If (a_1, a_2, a_3, a_4) is in $\text{supp}(f)$ with $a_4 > 0$ for $f \in K(\Pi) \cap K[\mathbf{x}]$, then $a_1 + a_2 + a_3$ is positive.*

Proof. Suppose to the contrary that $(0, 0, 0, a_4)$ is in $\text{supp}(f)$ with $a_4 > 0$ for some $f \in K(\Pi) \cap K[\mathbf{x}]$. Then, $\Phi^{-1}(f)$ is in $K[\mathbf{y}]^E$, since $K(\Pi) \cap K[\mathbf{x}] = K(\Pi) \cap K[\mathbf{x}]$ by Lemma 2.2 and $\Phi^{-1}(K(\Pi)) = K[\mathbf{y}]^E$. Note that any monomial in $K[\mathbf{y}] \setminus K[y_4]$ is not sent to $x_4^{a_4}$ by Φ . Hence, $l = a_4/\gamma$ is an integer, and y_4^l appears in $\Phi^{-1}(f)$. We remark that $E(\mathbf{y}^b) = \sum_{i=1}^4 b_i \mathbf{y}^b y_i^{-1}$ for $b = (b_1, b_2, b_3, b_4) \in (\mathbb{Z}_{\geq 0})^4$. Hence, for any $g \in K[\mathbf{y}]^E$ and $b \in \text{supp}(g)$, if a monomial \mathbf{y}^c appears in $E(\mathbf{y}^b)$ with nonzero coefficient, then there exists $d \in \text{supp}(g) \setminus \{b\}$ such that $\mathbf{y}^c = \mathbf{y}^d y_i^{-1}$ for some i . Actually, if such d did not exist, then \mathbf{y}^c would appear in $E(g)$. So, there appears in $\Phi^{-1}(f)$ a monomial $\mathbf{y}^d \neq y_4^l$ such that $y_4^{l-1} = \mathbf{y}^d y_i^{-1}$ for some i , since $E(\Phi^{-1}(f)) = 0$ and $E(y_4^l) = l y_4^{l-1}$. Clearly, \mathbf{y}^d is equal to $y_1 y_4^{l-1}$ or $y_2 y_4^{l-1}$ or $y_3 y_4^{l-1}$. Hence, one of $\mathbf{x}^{\delta_1} x_4^{a_4-\gamma}$, $\mathbf{x}^{\delta_2} x_4^{a_4-\gamma}$, $\mathbf{x}^{\delta_3} x_4^{a_4-\gamma}$ appears in f . This is a contradiction, since these monomials are not in $K[\mathbf{x}]$. \square

We define linear functions $l_i : \mathbf{R}^4 \rightarrow \mathbf{R}$ for $i = 1, 2, 3$ by

$$l_i((a_1, a_2, a_3, a_4)) = a_j + a_k - (a_1 + a_2 + a_3)\xi_i, \quad (2.3)$$

where $j, k \in \{1, 2, 3\} \setminus \{i\}$ with $j \neq k$. Let \mathcal{C} be the set of $a \in \mathbf{R}^4$ such that $l_i(a) \geq 0$ for $i = 1, 2, 3$. Then, \mathcal{C} is a convex polyhedral cone in \mathbf{R}^4 . We remark that $\text{supp}(f) \subset \mathcal{C}$ implies $\Phi(f) \in K[\mathbf{x}]$ for each $f \in K[\mathbf{y}]$. To verify this, we may assume that $f = \mathbf{y}^b$ for some $b = (b_1, b_2, b_3, b_4) \in (\mathbb{Z}_{\geq 0})^4$. Since $b \in \mathcal{C}$, we have

$$\begin{aligned} 0 &\leq (\delta_{i,i} + \min\{\delta_{j,i}, \delta_{k,i}\})l_i(b) \\ &= (b_j + b_k)(\delta_{i,i} + \min\{\delta_{j,i}, \delta_{k,i}\}) - (b_1 + b_2 + b_3)\delta_{i,i} \\ &= (b_j + b_k) \min\{\delta_{j,i}, \delta_{k,i}\} - b_i \delta_{i,i} \end{aligned} \quad (2.4)$$

for $i = 1, 2, 3$. The i th component of $a = \sum_{i=1}^3 b_i \delta_i + b_4 \gamma \mathbf{e}_4$ is not less than the right-hand side of the last equality in (2.4) for $i = 1, 2, 3$. The fourth component of a is always nonnegative. Thus, $\Phi(\mathbf{y}^b) = \mathbf{x}^a$ is in $K[\mathbf{x}]$.

The following is the key lemma.

Lemma 2.4. *There exist positive integers p_1, p_2 , and p_3 with the following properties:*

- (i) *The first three components of $p_3 \delta_2 + (p_1 + p_2) \delta_3$ are positive.*
- (ii) *For each positive integer q , there exists an element of $K[\mathbf{y}]^E$ of the form*

$$(y_3 - y_2)^{p_1} (y_3 - y_1)^{p_2} (y_2 - y_1)^{p_3} y_4^q + (\text{terms of lower degree in } y_4) \quad (2.5)$$

whose support is contained in \mathcal{C} .

We show Theorem 1.1 as a consequence of Lemma 2.4. Let S be the union of $\text{supp}(f)$ for $f \in K(\Pi) \cap K[\mathbf{x}]$. Then, the function $N : S \setminus \{0\} \rightarrow \mathbf{R}$ sending $a = (a_1, a_2, a_3, a_4) \in S \setminus \{0\}$ to $a_4/(a_1 + a_2 + a_3)$ is well-defined, since $a_1 + a_2 + a_3 = 0$ implies $a = 0$ by Lemma 2.3. Suppose to the contrary that $K(\Pi) \cap K[\mathbf{x}]$ is generated by a finite number of elements g_1, \dots, g_r . Then, S is contained in the subsemigroup of \mathbf{Z}^4 generated by $S' = \bigcup_{i=1}^r \text{supp}(g_i)$. Since $S' \setminus \{0\}$ is a finite subset of $S \setminus \{0\}$, there exists $\mu > 0$ such that $N(a) < \mu$ for every $a \in S' \setminus \{0\}$. Note that $N(a + b) \leq \max\{N(a), N(b)\}$ for a and b . Thus, we have $N(a) < \mu$ for any $a \in S \setminus \{0\}$. Take $p_1, p_2, p_3 > 0$ as in Lemma 2.4. Then, there exists an element of $K[\mathbf{y}]^E$ as in (2.5) whose support is contained in \mathcal{C} for each $q > 0$ by Lemma 2.4. It is sent to $f \in K(\Pi) \cap K[\mathbf{x}]$ of the form $f = f' + f''$ by Φ . Here,

$$f' = (\mathbf{x}^{\delta_3} - \mathbf{x}^{\delta_2})^{p_1} (\mathbf{x}^{\delta_3} - \mathbf{x}^{\delta_1})^{p_2} (\mathbf{x}^{\delta_2} - \mathbf{x}^{\delta_1})^{p_3} x_4^{q\gamma}$$

and f'' is an element of $K[\mathbf{x}]$ with $\text{supp}(f') \cap \text{supp}(f'') = \emptyset$. Hence, $c = p_3\delta_2 + (p_1 + p_2)\delta_3 + q\gamma\mathbf{e}_4$ is in S for any $q > 0$. This is a contradiction, since $N(c) > \mu$ for sufficiently large q . Therefore, $K(\Pi) \cap K[\mathbf{x}]$ is not finitely generated.

Note that c is a vertex of the Newton polytope of f , i.e., the convex hull of $\text{supp}(f)$ in \mathbf{R}^4 . Moreover, every component of c is not zero by Lemma 2.4. Let D be a nonzero locally nilpotent derivation on $K[\mathbf{x}]$. Then, by Hadas and Makar-Limanov's theorem [5, Theorem 3.2], at least one component of each vertex of the Newton polytope of an element of $K[\mathbf{x}]^D \setminus \{0\}$ must be zero. Hence, $f \notin K[\mathbf{x}]^D$. Therefore, $K(\Pi) \cap K[\mathbf{x}] \not\subset K[\mathbf{x}]^D$ for any nonzero locally nilpotent derivation D on $K[\mathbf{x}]$. Thus, the proof of Theorem 1.1 is completed on the assumption that Lemma 2.4 is true.

3. Proof of Lemma 2.4

In this section, we prove Lemma 2.4. For each $r \in \mathbf{R}$, we denote by $\lfloor r \rfloor$ the minimal integer m such that $m \geq r$. Note that $r \leq \lfloor r \rfloor < r + 1$ for $r \in \mathbf{R}$.

Lemma 3.1. *There exists a positive integer p_0 with the following property. For each integer $p \geq p_0$, there exist positive integers p_1 and p_2 with $p_1 \geq p\xi_1$ such that*

$$\begin{aligned} b = & (p - p_2 + r)\mathbf{e}_2 + (p_2 - r)\mathbf{e}_3 + \alpha_1 p'_1 (\mathbf{e}_3 - \mathbf{e}_2) + \alpha_2 (p_2 - r)(\mathbf{e}_1 - \mathbf{e}_3) \\ & + \alpha_3 (p - p'_1 - p_2 + r)(\mathbf{e}_1 - \mathbf{e}_2) \end{aligned} \quad (3.1)$$

satisfies $l_2(b), l_3(b) \geq 0$ for any $\alpha_1, \alpha_2, \alpha_3 \in [0, 1]$, $0 \leq p'_1 \leq p_1$, and $r = 0$. If $p'_1 = p_1$, then b is contained in \mathcal{C} for any $\alpha_1, \alpha_2, \alpha_3 \in [0, 1]$ and $r \in \{-1, 0, 1\}$.

Proof. By hypothesis, $\xi_1 + \xi_2 + \xi_3 < 1$. Let p_0 be a positive integer such that $p_0(1 - \xi_1 - \xi_2 - \xi_3) \geq 4$, and p an integer such that $p \geq p_0$. Set $p_1 = p - \lfloor p\xi_2 \rfloor - \lfloor p\xi_3 \rfloor - 2$ and $p_2 = \lfloor p\xi_2 \rfloor + 1$. Then, we have $p_1 > p(1 - \xi_2 - \xi_3) - 4 \geq p\xi_1$. For $r \in \{-1, 0, 1\}$ and $0 \leq p'_1 \leq p_1$, put

$$\begin{aligned}
b_1 &= (p - p_2 + r)\mathbf{e}_2 + (p_2 - r)\mathbf{e}_3, & b_2 &= b_1 + (p_2 - r)(\mathbf{e}_1 - \mathbf{e}_3), \\
b_3 &= b_2 + (p - p'_1 - p_2 + r)(\mathbf{e}_1 - \mathbf{e}_2), & b_4 &= b_1 + p'_1(\mathbf{e}_3 - \mathbf{e}_2), \\
b_5 &= b_4 + (p - p'_1 - p_2 + r)(\mathbf{e}_1 - \mathbf{e}_2), & b_6 &= b_5 + (p_2 - r)(\mathbf{e}_1 - \mathbf{e}_3).
\end{aligned}$$

Then, b in (3.1) is contained in the convex hull of b_1, \dots, b_6 in \mathbf{R}^4 for each $\alpha_1, \alpha_2, \alpha_3 \in [0, 1]$. Hence, to complete the proof, it suffices to verify for $j = 1, \dots, 6$ and $r \in \{-1, 0, 1\}$ that $l_2(b_j), l_3(b_j) \geq 0$ for $0 \leq p'_1 \leq p_1$ and $l_1(b_j) \geq 0$ for $p'_1 = p_1$.

For $0 \leq p'_1 \leq p_1$ and $r \in \{-1, 0, 1\}$, we have

$$\begin{aligned}
l_2(b_1) &= p_2 - r - p\xi_2 = \lfloor p\xi_2 \rfloor + 1 - r - p\xi_2 \geq p\xi_2 + 1 - r - p\xi_2 = 1 - r \geq 0, \\
l_3(b_1) &= p - (\lfloor p\xi_2 \rfloor + 1) + r - p\xi_3 > p(1 - \xi_2 - \xi_3) + r - 2 \geq p\xi_1 + r + 2 > 0. \quad (3.2)
\end{aligned}$$

Note that $p - p'_1 - p_2 + r \geq p - p_1 - p_2 + r = \lfloor p\xi_3 \rfloor + 1 + r > 0$ and $p_2 - r = \lfloor p\xi_2 \rfloor - r + 1 > 0$. Moreover,

$$l_i(\mathbf{e}_3 - \mathbf{e}_2) = \begin{cases} 0 & \text{if } i = 1, \\ 1 & \text{if } i = 2, \\ -1 & \text{if } i = 3, \end{cases} \quad \text{and} \quad l_j(\mathbf{e}_1 - \mathbf{e}_k) = \begin{cases} -1 & \text{if } j = 1, k \in \{2, 3\}, \\ 1 & \text{if } j, k \in \{2, 3\}, j = k, \\ 0 & \text{if } j, k \in \{2, 3\}, j \neq k. \end{cases}$$

Hence, $l_2(b_1) \leq l_2(b_4)$ and $l_i(b_1) \leq l_i(b_2) \leq l_i(b_3)$, $l_i(b_4) \leq l_i(b_5) \leq l_i(b_6)$ for $i = 2, 3$. Thus, we get $l_2(b_j), l_3(b_j) \geq 0$ for each j for $0 \leq p'_1 \leq p_1$ by (3.2) and

$$l_3(b_4) = l_3(b_1) + p'_1 l_3(\mathbf{e}_3 - \mathbf{e}_2) = l_3(b_1) - p'_1 \geq l_3(b_1) - p_1 \geq r + 1 \geq 0.$$

Since $l_1(b_1) = l_1(b_4)$, we have $l_1(b_3) = l_1(b_6)$. Furthermore, $l_1(b_1) > l_1(b_2) > l_1(b_3)$ and $l_1(b_4) > l_1(b_5) > l_1(b_6)$. If $p'_1 = p_1$, then

$$l_1(b_3) = l_1(b_1) - (p_2 - r) - (p - p_1 - p_2 + r) = -p\xi_1 + p_1 > 0.$$

Therefore, we get $l_1(b_j) \geq 0$ for each j if $p'_1 = p_1$. \square

Now, take p_0 as in Lemma 3.1, and choose p_1 and p_2 for $p = p_0$. We set

$$\begin{aligned}
f_r &= (y_3 - y_2)^{p_1} (y_3 - y_1)^{p_2 - r} (y_2 - y_1)^{p_3 + r} \\
&= y_2^{p - p_2 + r} y_3^{p_2 - r} (y_2^{-1} y_3 - 1)^{p_1} (1 - y_1 y_3^{-1})^{p_2 - r} (1 - y_1 y_2^{-1})^{p_3 + r}
\end{aligned}$$

for $r \in \{-1, 0, 1\}$, where $p_3 = p - p_1 - p_2$. Then, each element of $\text{supp}(f_r)$ is written as b in (3.1) with $p'_1 = p_1$ for some $\alpha_1, \alpha_2, \alpha_3 \in [0, 1]$. Hence, $\text{supp}(f_r)$ is contained in \mathcal{C} by Lemma 3.1, and so $\Phi(f_r)$ is in $K[\mathbf{x}]$ for each r . In particular, $(p_3 + r)\delta_2 + (p_1 + p_2 - r)\delta_3$ is in $(\mathbf{Z}_{\geq 0})^4$ for each r . This implies that the first three components of $p_3\delta_2 + (p_1 + p_2)\delta_3$ are positive. Thus, the first statement of Lemma 2.4 is proved.

We remark that, for $f' \in K[\mathbf{y}]$ and $e' \in \mathbf{Z}_{\geq 0}$, if $l_2(b), l_3(b) \geq 0$ for any $b \in \text{supp}(f')$, then each of $b \in \text{supp}(f'(y_4 - y_1)^{e'})$ also satisfies $l_2(b), l_3(b) \geq 0$. Actually, we have $l_i(\mathbf{e}_4), l_i(\mathbf{e}_1 - \mathbf{e}_4) \geq 0$ for $i = 2, 3$. Take any positive integer q , and let

\mathcal{F} be the set of homogeneous $(p+q)$ -forms $F \in K[\mathbf{y}]^E$ of the form $F = f_0 y_4^q +$ (terms of lower degree in y_4) such that $l_2(b), l_3(b) \geq 0$ for any $b \in \text{supp}(F)$. Since $\text{supp}(f_0) \subset \mathcal{C}$, we have $f_0(y_4 - y_1)^q \in \mathcal{F}$, as remarked. So, \mathcal{F} is not empty. To complete the proof of the second statement of Lemma 2.4, it suffices to show the existence of $F \in \mathcal{F}$ such that $l_1(b) \geq 0$ for any $b \in \text{supp}(F)$. Suppose the contrary, that is, there exists $b \in \text{supp}(F)$ such that $l_1(b) < 0$ for every $F \in \mathcal{F}$. We define $O(F) = (d, e) \in \mathbf{Z}^2$ for each $F \in \mathcal{F}$ by

$$\begin{aligned} e &= \max\{b_4 \mid b = (b_1, b_2, b_3, b_4) \in \text{supp}(F), l_1(b) < 0\}, \\ d &= \max\{b_1 \mid b = (b_1, b_2, b_3, b_4) \in \text{supp}(F), b_4 = e\}. \end{aligned}$$

Consider the total order \preccurlyeq on \mathbf{Z}^2 defined by $(d_1, e_1) \preccurlyeq (d_2, e_2)$ if $e_1 < e_2$ or $e_1 = e_2$, $d_1 \leq d_2$. Then, choose $F \in \mathcal{F}$ such that $O(F) \preccurlyeq O(F')$ for any $F' \in \mathcal{F}$, and let $h \in K[y_2, y_3]$ be the coefficient of $y_1^d y_4^e$ in F , where $O(F) = (d, e)$. Put $s = p + q - d - e$. If b is an element of $\text{supp}(F)$ with first and fourth components d and e , respectively, then $l_1(b) = s - (p + q - e)\xi_1 = (p + q - e)(1 - \xi_1) - d$. By the maximality of d , the right-hand side of this equality is negative. Hence, we get

$$s < (p + q - e)\xi_1. \quad (3.3)$$

Lemma 3.2. *There exists $\beta \in K \setminus \{0\}$ such that $h = \beta(y_3 - y_2)^s$.*

Proof. It suffices to show that $E(h) = 0$. Indeed, $K[y_2, y_3]^E = K[y_3 - y_2]$, and h is a homogeneous s -form in y_2 and y_3 . Suppose to the contrary that a monomial $y_2^{b_2} y_3^{b_3}$ appears in $E(h)$ with nonzero coefficient. Then, $c - \mathbf{e}_i = (d, b_2, b_3, e)$ for some $c \in \text{supp}(F)$ and $i \in \{1, 4\}$, since $E(F) = 0$. By the maximality of d , we have $i \neq 1$. Hence, $c = (d, b_2, b_3, e + 1)$. Since $b_2 + b_3 = p + q - d - e - 1$ and $\xi_1 < 1$, we have

$$l_1(c) = b_2 + b_3 - \xi_1(d + b_2 + b_3) = s - (p + q - e)\xi_1 + (\xi_1 - 1) < 0$$

by (3.3). This contradicts the maximality of e . Therefore, $E(h) = 0$. \square

We put $\tilde{p} = p + q - e$. Then, $\tilde{p} > p$, since $0 \leq e < q$. So, by Lemma 3.1, there exist positive integers \tilde{p}_1 and \tilde{p}_2 with $\tilde{p}_1 \geq \tilde{p}\xi_1$ such that

$$\begin{aligned} b &= (\tilde{p} - \tilde{p}_2)\mathbf{e}_2 + \tilde{p}_2\mathbf{e}_3 + \alpha_1 p'_1(\mathbf{e}_3 - \mathbf{e}_2) + \alpha_2 \tilde{p}_2(\mathbf{e}_1 - \mathbf{e}_3) \\ &\quad + \alpha_3(\tilde{p} - p'_1 - \tilde{p}_2)(\mathbf{e}_1 - \mathbf{e}_2) \end{aligned} \quad (3.4)$$

satisfies $l_2(b), l_3(b) \geq 0$ for any $\alpha_1, \alpha_2, \alpha_3 \in [0, 1]$ and $0 \leq p'_1 \leq \tilde{p}_1$. Every element of the support of $g = (y_3 - y_2)^s (y_1 - y_3)^{\tilde{p}_2} (y_1 - y_2)^{\tilde{p} - s - \tilde{p}_2}$ is expressed as b in (3.4) with $p'_1 = s$ for some $\alpha_1, \alpha_2, \alpha_3 \in [0, 1]$. Since $0 < s < \tilde{p}\xi_1 \leq \tilde{p}_1$ by (3.3), we get $l_2(b), l_3(b) \geq 0$ for any $b \in \text{supp}(g)$. Hence, $l_2(b), l_3(b) \geq 0$ for any $b \in \text{supp}(g(y_4 - y_1)^e)$, as remarked above. So, $H = F - \beta g(y_4 - y_1)^e$ is in \mathcal{F} . Note that $g(y_4 - y_1)^e$ is written as $(y_3 - y_2)^s y_1^d y_4^e + w$, where $w \in K[\mathbf{y}]$ such that the first and the fourth component d' and e' of each $b \in \text{supp}(w)$ satisfy $(d', e') \preccurlyeq (d, e)$ and $(d', e') \neq (d, e)$. Since $h = \beta(y_3 - y_2)^s$ by Lemma 3.2, we

have $O(H) \preceq O(F)$ and $O(H) \neq O(F)$. This contradicts the choice of F . Therefore, the proof of Lemma 2.4 is completed.

Acknowledgments

The author thanks to Professor Masanori Ishida for kind encouragement, to Professors Harm Derksen and Gene Freudenburg for informing him of the Fourteenth Problem of Hilbert in dimension three, and to Dr. Ryuji Tanimoto for reading the first version of this paper and giving him valuable comments. He also thanks the referee for pointing out that our new counterexample is not equal to any invariant subring of $K[\mathbf{x}]$ as mentioned in Section 1.

The final version of this paper was prepared when the author visited the Department of Mathematics of the University of Michigan. He is grateful to his host Professor Karen Smith.

References

- [1] D. Daigle, G. Freudenburg, A counterexample to Hilbert's fourteenth problem in dimension 5, *J. Algebra* 221 (1999) 528–535.
- [2] D. Daigle, G. Freudenburg, Triangular derivations of $\mathbf{k}[X_1, X_2, X_3, X_4]$, *J. Algebra* 241 (2001) 328–339.
- [3] A. van den Essen, Polynomial automorphisms and the Jacobian conjecture, in: *Progr. Math.*, vol. 190, Birkhäuser, Basel, 2000.
- [4] G. Freudenburg, A counterexample to Hilbert's fourteenth problem in dimension six, *Transform. Groups* 5 (2000) 61–71.
- [5] O. Hadas, L. Makar-Limanov, Newton polytopes of constants of locally nilpotent derivations, *Comm. Algebra* 28 (2000) 3667–3678.
- [6] H. Kojima, M. Miyanishi, On Roberts' counterexample to the fourteenth problem of Hilbert, *J. Pure Appl. Algebra* 122 (1997) 277–292.
- [7] S. Kuroda, A condition for finite generation of the kernel of a derivation, *J. Algebra* 262 (2003) 391–400.
- [8] S. Kuroda, A generalization of Roberts' counterexample to the fourteenth problem of Hilbert, *Tohoku Math. J.*, in press.
- [9] S. Kuroda, A finite universal SAGBI basis for the kernel of a derivation, *Osaka J. Math.*, in press.
- [10] S. Kuroda, A counterexample to the fourteenth problem of Hilbert in dimension three, *Michigan Math. J.*, in press.
- [11] S. Maubach, Triangular monomial derivations on $k[X_1, X_2, X_3, X_4]$ have kernel generated by at most four elements, *J. Pure Appl. Algebra* 153 (2000) 165–170.
- [12] M. Nagata, On the fourteenth problem of Hilbert, in: *Proc. of the Internat. Congr. of Mathematicians*, 1958, Cambridge Univ. Press, London, 1960, pp. 459–462.
- [13] M. Nagata, On the 14th problem of Hilbert, *Amer. J. Math.* 81 (1959) 766–772.
- [14] P. Roberts, An infinitely generated symbolic blow-up in a power series ring and a new counterexample to Hilbert's fourteenth problem, *J. Algebra* 132 (1990) 461–473.
- [15] O. Zariski, Interprétations algébrique-géométriques du quatorzième problème de Hilbert, *Bull. Sci. Math.* 78 (1954) 155–168.